

tdrd1^{fh244}

Nature of the mutation

The *fh244* allele contains a single G-to-T point mutation that results in a premature stop codon (Moens C., personal communication).

Genotyping assay

Genotyping of the *fh244* allele is based on the RFLP assay (**R**estriction **F**ragment **L**ength **P**olymorphism; Botstein *et al.*, Am. J. Hum. Genet. 32: 314-331, 1980). This method is used to detect a mutation that either creates or abolishes a site recognized by a specific restriction enzyme. In the RFLP assay, a sequence of interest is first PCR-amplified and then the PCR product is subjected to restriction enzyme digestion. The presence or absence of the mutation is determined by the resulting restriction pattern. The *fh244* mutation abolishes a site recognized by the Hpy188I restriction enzyme.

Primers:

fh244_03: 5' GAA AAA CCT AAG GAG TCA AAA GCT G 3'

fh244_04: 5' GGC AGA GTG TCT ATG CTT GGA TAA C 3'

PCR program (55_30_30):

1. 94°C for 3 min
2. 94°C for 30 sec
3. **55°C for 30 sec**
4. 72°C for **30 sec**
5. Go to step 2 (above) for 39 cycles
6. 72°C for 5 min
7. 8.0°C hold
8. END

Product size: 277 bp

Digestion of the PCR product with the Hpy188I restriction enzyme:

Product type	Product digestion	DNA fragments after digestion (bp)
PCR product derived from the WT template	cleaved	210 bp and 67 bp
PCR product containing the mutation	unaffected	277 bp

IMPORTANT NOTE: It is highly recommended to use WT positive controls to monitor whether enzyme digestion has been carried out to completion. Without this control, partially digested WT samples can be mistakenly regarded as heterozygous samples.

Zebrafish International Resource Center (ZIRC)

5274 University of Oregon
Eugene, OR 97403-5274, USA
Phone: 541-346-6028
Email: genotyping@zebrafish.org
Web: <http://zebrafish.org>