



ift88^{tz288}

Nature of the mutation

The *tz288* allele contains a single T-to-A point mutation that introduces a premature stop codon in exon 11 of the *ift88* transcription unit (Tsujikawa and Malicki, Neuron 42(5): 703-716, 2004).

Genotyping assay

Genotyping of the *tz288* allele is based on the RFLP assay (**R**estriction **F**ragment **L**ength **P**olymorphism; Botstein *et al.*, Am. J. Hum. Genet. 32: 314-331, 1980). This method is used to detect a mutation that either creates or abolishes a site recognized by a specific restriction enzyme. In the RFLP assay, a sequence of interest is first PCR-amplified and then the PCR product is subjected to restriction enzyme digestion. The presence or absence of the mutation is determined by the resulting restriction pattern. The *tz288* mutation creates a site recognized by the DdeI restriction enzyme.

Primers:

OVL_01: 5' GCA AAT TAG TGC ATA TAA CGC CTC 3'

OVL_02: 5' CCA ATG TTC TGC ATG ATC TTT ATC C 3'

PCR program (53_30_30):

1. 94°C for 3 min
2. 94°C for 30 sec
3. **53°C for 30 sec**
4. 72°C for **30 sec**
5. Go to step 2 (above) for 39 cycles
6. 72°C for 5 min
7. 8.0°C hold
8. END

Product size: 401 bp

Digestion of the PCR product with the DdeI restriction enzyme:

| Product type | Product digestion | DNA fragments after digestion (bp) |
|--|-------------------|------------------------------------|
| PCR product derived from the WT template | unaffected | 401 bp |
| PCR product containing the mutation | cleaved | 259 bp and 142 bp |

Zebrafish International Resource Center (ZIRC)

5274 University of Oregon
Eugene, OR 97403-5274, USA
Phone: 541-346-6028
Email: genotyping@zebrafish.org
Web: <http://zebrafish.org>