

Tg(UAS:myc-Notch1a-intra)kca3

Transgene description

The *Tg(UAS:myc-Notch1a-intra)kca3* transgene contains the *myc-Notch1a-intra* chimeric cDNA under control of the UAS promoter (Scheer and Campos-Ortega, *Mechanisms of Development* 80:153-158, 1999).

Genotyping assay

To genotype the *Tg(UAS:myc-Notch1a-intra)kca3* line, the transgene-specific primers (**P009.F** and **P010.R**) are used.

Primers:

P009.F: 5' CAT CGC GTC TCA GCC TCA C 3'

P010.R: 5' CGG AAT CGT TTA TTG GTG TCG 3'

PCR program (61_30_45):

- 1_94°C for 3 min
- 2_94°C for 30 s
- 3_61°C for 30 s
- 4_72°C for 45 s
- 5_Go to step 2 (above) for 34 cycles
- 6_72°C for 5 min
- 7_8.0°C hold
- 8_END

Product size: 450 bp

The 450-bp product is specific for the genomic DNA containing the *Tg(UAS:myc-Notch1a-intra)kca3* transgene. No PCR product is generated for wild-type genomic DNA.

IMPORTANT NOTE: It is possible that multiple copies of the transgene might have integrated into the genome during transgenesis and that some of these integrations are non-functional. Samples that contain only a non-functional transgene or its fragment will be identified falsely as positive in the genotyping assay. For this reason, it is recommended to use functional assays to verify individuals identified as positive in the genotyping assay.

Zebrafish International Resource Center (ZIRC)

5274 University of Oregon
Eugene, OR 97403-5274, USA
Phone: 541-346-6028
Email: genotyping@zebrafish.org
Web: <http://zebrafish.org>