

Tg(hsp70l:tcf3-GFP)w26

Transgene description

The *Tg(hsp70l:tcf3-GFP)w26* transgene contains the *tcf3-GFP* chimeric cDNA under control of the *hsp70l* promoter (Dorsky R., personal communication).

Genotyping assay

To genotype the *Tg(hsp70l:tcf3-GFP)w26* line, the transgene-specific primers (**GFP003** and **GFP004**) are used.

Primers:

GFP003: 5' AGA GGG TGA AGG TGA TGC AAC ATA C 3'

GFP004: 5' TGG TCT GCT AGT TGA ACG CTT CC 3'

PCR program (55_30_30):

- 1_94°C for 3 min
- 2_94°C for 30 s
- 3_55°C for 30 s
- 4_72°C for 30 s
- 5_Go to step 2 (above) for 34 cycles
- 6_72°C for 5 min
- 7_8.0°C hold
- 8_END

Product size: 450 bp

The 450-bp product is specific for the genomic DNA containing the *Tg(hsp70l:tcf3-GFP)w26* transgene. No PCR product is generated for wild-type genomic DNA.

IMPORTANT NOTE: It is possible that multiple copies of the transgene might have integrated into the genome during transgenesis and that some of these integrations are non-functional. Samples that contain only a non-functional transgene or its fragment will be identified falsely as positive in the genotyping assay. For this reason, it is recommended to use functional assays to verify individuals identified as positive in the genotyping assay.